

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant SOX10 Antibody, Clone: [SBX10-2R], Rabbit, Monoclonal NSJ-V3777IHC-7ML

| | |
|--------------------------|---|
| Artikelname | Recombinant SOX10 Antibody, Clone: [SBX10-2R], Rabbit, Monoclonal |
| Artikelnummer | NSJ-V3777IHC-7ML |
| Hersteller Artikelnummer | V3777IHC-7ML |
| Alternativnummer | NSJ-V3777IHC-7ML |
| Hersteller | NSJ Bioreagents |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | Amino acids 115-269 of the human protein were used as the immunogen for the recombinant SOX10 antibody. |
| Produktbeschreibung | SOX10 is a member of the SOX (SRV-related HMG-box) family of transcription factors involved in the regulation of embryonic development and in the determination of the cell fate. It may act as a transcriptional activator after forming a protein comple... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [SBX10-2R] |
| UniProt | P56693 |
| Reinheit | Protein A affinity chromatography |

| | |
|------------------------|---|
| Formulierung | Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only* |
| Antibody Type | Primary Antibody |
| Application Verdünnung | The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |
| Anwendungsbeschreibung | Titering of the recombinant SOX10 antibody may be required for optimal performance.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |