

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD22 Antibody, Clone: [BLCAM/1796], Mouse, Monoclonal NSJ-V3784-100UG

| | |
|--------------------------|---|
| Artikelname | CD22 Antibody, Clone: [BLCAM/1796], Mouse, Monoclonal |
| Artikelnummer | NSJ-V3784-100UG |
| Hersteller Artikelnummer | V3784-100UG |
| Alternativnummer | NSJ-V3784-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | ELISA, FACS, IF, WB |
| Spezies Reaktivität | Human |
| Immunogen | Amino acids 52-178 from the human protein were used as the immunogen for the CD22 antibody. |
| Produktbeschreibung | Recognizes a protein of 130-140kDa, identified as CD22 (also known as BL-CAM). CD22 expression is restricted to normal and neoplastic B cells and is absent from other haemopoietic cell types. In B-cell ontogeny, CD22 is first expressed in the cytopla... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [BLCAM/1796] |
| UniProt | P20273 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|---|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | ELISA: 2-4ug/ml (order BSA/azide-free format),Western blot: 1-2ug/ml,Flow cytometry: 1-2ug/10 ⁶ cells,Immunofluorescence: 1-2ug/ml |
| Anwendungsbeschreibung | Optimal dilution of the CD22 antibody should be determined by the researcher. |