

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **NR3C2 Antibody / Mineralocorticoid Receptor, Clone: [PCRP-NR3C2-1E3], Mouse, Monoclonal NSJ-V5075-20UG**

|                          |  |
|--------------------------|--|
| Artikelname              | NR3C2 Antibody / Mineralocorticoid Receptor, Clone: [PCRP-NR3C2-1E3], Mouse, Monoclonal  |
| Artikelnummer            | NSJ-V5075-20UG   |
| Hersteller Artikelnummer | V5075-20UG   |
| Alternativnummer         | NSJ-V5075-20UG   |
| Hersteller               | NSJ Bioreagents  |
| Wirt                     | Mouse  |
| Kategorie                | Antikörper   |
| Applikation              | FACS   |
| Spezies Reaktivität      | Human  |
| Immunogen                | A recombinant partial protein sequence (within amino acids 601-673) from the human protein was used as the immunogen for the NR3C2 antibody.   |
| Produktbeschreibung      | Mineralocorticoid hormones are primarily found in epithelial tissues where they function as regulators of Na <sup>+</sup> , K <sup>+</sup> and H <sup>+</sup> ion transport. Aldosterone is a mineralocorticoid that has been shown to regulate electrolyte excretion and intravascular volume ... |
| Klonalität               | Monoclonal   |
| Klon-Bezeichnung         | [PCRP-NR3C2-1E3]   |
| UniProt                  | <a href="#">P08235</a>   |

|                        |  |
|------------------------|--|
| Reinheit               | Protein A/G affinity   |
| Formulierung           | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide        |
| Antibody Type          | Primary Antibody   |
| Application Verdünnung | Flow cytometry: 1-2ug/million cells  |
| Anwendungsbeschreibung | Optimal dilution of the NR3C2 antibody should be determined by the researcher. |