

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

ICOS Ligand Antibody / ICOS-L / B7RP-1, Clone: [ISLG-1], Mouse, Monoclonal NSJ-V7532-100UG

Artikelname	ICOS Ligand Antibody / ICOS-L / B7RP-1, Clone: [ISLG-1], Mouse, Monoclonal
Artikelnummer	NSJ-V7532-100UG
Hersteller Artikelnummer	V7532-100UG
Alternativnummer	NSJ-V7532-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	FACS, IF, IHC-P
Spezies Reaktivität	Human
Immunogen	A recombinant fragment of human ICOSLG / B7RP-1 protein within amino acids 23-149 was used as the immunogen for the ICOS Ligand antibody.
Produktbeschreibung	Ligand for the T-cell-specific cell surface receptor ICOS. Acts as a costimulatory signal for T-cell proliferation and cytokine secretion, induces also B-cell proliferation and differentiation into plasma cells. Could play an important role in mediat...
Klonalität	Monoclonal
Klon-Bezeichnung	[ISLG-1]
UniProt	O75144

Reinheit	Protein G affinity chromatography
Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Flow cytometry: 1-2ug/million cells,Immunofluorescence: 1-2ug/ml,Immunohistochemistry (FFPE): 1-2ug/ml for 30 minutes at RT,Prediluted IHC only format: incubate for 30 min at RT (1)
Anwendungsbeschreibung	Optimal dilution of the ICOS Ligand antibody should be determined by the researcher.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.