

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant ZAP70 Antibody, Clone: [ZAP70/4040R], Rabbit, Monoclonal NSJ-V8750-20UG

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | Recombinant ZAP70 Antibody, Clone: [ZAP70/4040R], Rabbit, Monoclonal |
| Artikelnummer | NSJ-V8750-20UG |
| Hersteller Artikelnummer | V8750-20UG |
| Alternativnummer | NSJ-V8750-20UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | ELISA, IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | A portion of amion acids 200-350 from the human protein was used as the immunogen for the recombinant ZAP70 antibody. |
| Produktbeschreibung | Zeta-associated protein-70 (ZAP-70) is a member of the Syk family of tyrosine kinases, a group of proteins that attach to the zeta chain components of T-cell receptors to signal downstream events involved in the regulation of cell function, prolifera... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [ZAP70/4040R] |
| UniProt | P43403 |
| Reinheit | Protein A affinity chromatography |

| | |
|------------------------|---------------------------------------------------------------------------------------------------------------|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | ELISA: 2-4ug/ml for coating (order Ab without BSA),Immunohistochemistry (FFPE): 1-2ug/ml for 30 minutes at RT |
| Anwendungsbeschreibung | Optimal dilution of the recombinant ZAP70 antibody should be determined by the researcher. |