

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p504S Antibody / AMACR, Clone: [13H4], Rabbit, Monoclonal NSJ-V8770-100UG

| | |
|--------------------------|---|
| Artikelname | p504S Antibody / AMACR, Clone: [13H4], Rabbit, Monoclonal |
| Artikelnummer | NSJ-V8770-100UG |
| Hersteller Artikelnummer | V8770-100UG |
| Alternativnummer | NSJ-V8770-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC-P, WB |
| Spezies Reaktivität | Human |
| Immunogen | Full length human recombinant protein was used as the immunogen for the p504S antibody. |
| Produktbeschreibung | This antibody recognizes a protein of 44kDa, which is identified as AMACR, also known as p504S. It is an enzyme that is involved in bile acid biosynthesis and beta-oxidation of branched-chain fatty acids. AMACR is essential in lipid metabolism. It is... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [13H4] |
| UniProt | Q9UHK6 |
| Reinheit | Protein A affinity chromatography |

| | |
|------------------------|---|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml,Western blot: 1-2ug/ml |
| Anwendungsbeschreibung | Optimal dilution of the p504S antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |