

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **hCG alpha Antibody / Luteinizing Hormone alpha, Clone: [SPM552], Mouse, Monoclonal NSJ-V9004-100UG**

|                          |                                                                                                                                                                                                                                                                        |
|--------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | hCG alpha Antibody / Luteinizing Hormone alpha, Clone: [SPM552],<br>Mouse, Monoclonal                                                                                                                                                                                  |
| Artikelnummer            | NSJ-V9004-100UG                                                                                                                                                                                                                                                        |
| Hersteller Artikelnummer | V9004-100UG                                                                                                                                                                                                                                                            |
| Alternativnummer         | NSJ-V9004-100UG                                                                                                                                                                                                                                                        |
| Hersteller               | NSJ Bioreagents                                                                                                                                                                                                                                                        |
| Wirt                     | Mouse                                                                                                                                                                                                                                                                  |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                             |
| Applikation              | IHC-P                                                                                                                                                                                                                                                                  |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                                  |
| Immunogen                | Recombinant hCG alpha protein was used as the immunogen for this<br>hCG alpha antibody.                                                                                                                                                                                |
| Produktbeschreibung      | HCG is a glycoprotein which is secreted in large quantities by normal<br>trophoblasts. It is present only in trace amounts in non-pregnant<br>urine and sera but rises sharply during pregnancy. HCG is composed<br>of two non-identical, non-covalently linked pol... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                             |
| Klon-Bezeichnung         | [SPM552]                                                                                                                                                                                                                                                               |
| UniProt                  | <a href="#">P01215</a>                                                                                                                                                                                                                                                 |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Reinheit               | Protein G affinity chromatography                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Formulierung           | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide                                                                                                                                                                                                                                                                                                                                                                                           |
| Antibody Type          | Primary Antibody                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT                                                                                                                                                                                                                                                                                                                                                                                                               |
| Anwendungsbeschreibung | <p>The optimal dilution of the hCG alpha antibody for each application should be determined by the researcher.1. No special pre-treatment is required for staining of formalin-fixed tissues. Note that HIER is NOT recommended for this antibody.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p> |