

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD35 Antibody (Complement Receptor 1), Clone: [SPM554], Mouse, Monoclonal NSJ-V9010-100UG

| | |
|--------------------------|---|
| Artikelname | CD35 Antibody (Complement Receptor 1), Clone: [SPM554], Mouse, Monoclonal |
| Artikelnummer | NSJ-V9010-100UG |
| Hersteller Artikelnummer | V9010-100UG |
| Alternativnummer | NSJ-V9010-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | Human monocytes were used as the immunogen for this CD35 antibody. |
| Produktbeschreibung | Recognizes a protein of 210-220kDa, which is identified as the Complement receptor 1 or CD35. This mAb is specific for a site in CR1 that is distal from the C3b-binding site, so that it is unable to block CR1/CD35 activity. It is highly specific and ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [SPM554] |
| UniProt | P17927 |

| | |
|------------------------|---|
| Reinheit | Protein G affinity chromatography |
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Anwendungsbeschreibung | <p>The optimal dilution of the CD35 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p> |