

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Glial Fibrillary Acidic Protein Antibody, Clone: [SPM248], Mouse, Monoclonal NSJ-V9022-100UG

Artikelname	Glial Fibrillary Acidic Protein Antibody, Clone: [SPM248], Mouse, Monoclonal
Artikelnummer	NSJ-V9022-100UG
Hersteller Artikelnummer	V9022-100UG
Alternativnummer	NSJ-V9022-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	FACS, IHC-P, WB
Spezies Reaktivität	Human
Immunogen	GFAP isolated from pig spinal cord was used as the immunogen for this Glial Fibrillary Acidic Protein antibody.
Produktbeschreibung	Glial Fibrillary Acidic Protein is specifically found in astroglia, making it a very popular marker for localizing benign astrocyte and neoplastic cells of glial origin in the central nervous system. Ab to GFAP is useful in differentiating primary gl...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM248]
UniProt	P14136

Reinheit	Protein G affinity chromatography
Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Flow cytometry: 1-2ug/10 ⁶ cells, Western blot: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT (1) (2)
Anwendungsbeschreibung	The optimal dilution of the Glial Fibrillary Acidic Protein antibody for each application should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.