

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

IgG Antibody, Clone: [SPM556], Mouse, Monoclonal NSJ-V9028IHC-7ML

Artikelname	IgG Antibody, Clone: [SPM556], Mouse, Monoclonal
Artikelnummer	NSJ-V9028IHC-7ML
Hersteller Artikelnummer	V9028IHC-7ML
Alternativnummer	NSJ-V9028IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Purified human Ig Gamma Chain was used as the immunogen for this IgG antibody.
Produktbeschreibung	Recognizes a protein of 75kDa, identified as gamma heavy chain of human immunoglobulins. It does not cross-react with IgA, IgM, IgE, or IgD, heavy chains, T-cells, monocytes, granulocytes, or erythrocytes. This mAb is useful in the identification of ...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM556]
UniProt	P01857
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	The optimal dilution of the IgG antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.