

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

IgM Antibody, Clone: [SPM557], Mouse, Monoclonal NSJ-V9029-20UG

| | |
|--------------------------|---|
| Artikelname | IgM Antibody, Clone: [SPM557], Mouse, Monoclonal |
| Artikelnummer | NSJ-V9029-20UG |
| Hersteller Artikelnummer | V9029-20UG |
| Alternativnummer | NSJ-V9029-20UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant heavy chain of human IgM was used as the immunogen for this IgM antibody. |
| Produktbeschreibung | Recognizes a protein of 75kDa, identified as mu heavy chain of human immunoglobulins. It does not cross-react with alpha (IgA), gamma (IgG), epsilon (IgE), or delta (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes. This mAb is u... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [SPM557] |
| UniProt | P01871 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|--|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Anwendungsbeschreibung | <p>The optimal dilution of the IgM antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p> |