

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Kappa Light Chain Antibody, Clone: [SPM508], Mouse, Monoclonal NSJ-V9030-20UG

Artikelname	Kappa Light Chain Antibody, Clone: [SPM508], Mouse, Monoclonal
Artikelnummer	NSJ-V9030-20UG
Hersteller Artikelnummer	V9030-20UG
Alternativnummer	NSJ-V9030-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Human B-lymphoma cells were used as the immunogen for this Kappa Light Chain antibody.
Produktbeschreibung	This mAb is specific to kappa light chain of immunoglobulin and shows no cross-reaction with lambda light chain or any of the five heavy chains. In mammals, the two light chains in an antibody are always identical, with only one type of light chain, ...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM508]
UniProt	<a href="#">P01601</a>
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	<p>The optimal dilution of the Kappa Light Chain antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p>