

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 18 Antibody, Clone: [SPM265], Mouse, Monoclonal NSJ-V9040-20UG

Artikelname	Cytokeratin 18 Antibody, Clone: [SPM265], Mouse, Monoclonal
Artikelnummer	NSJ-V9040-20UG
Hersteller Artikelnummer	V9040-20UG
Alternativnummer	NSJ-V9040-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IF, IHC-P, WB
Spezies Reaktivität	Human
Immunogen	The human breast cancer cell line PMC-42 was used as the immunogen for this Cytokeratin 18 antibody.
Produktbeschreibung	This mAb reacts with a wide variety of simple epithelia. It does not react with stratified squamous epithelia. It reacts with epithelial tumors of the gastrointestinal tract, lung, breast, pancreas, ovary, and thyroid. Cytokeratin 18, which belongs t...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM265]
UniProt	P05783
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunofluorescence: 1-2ug/ml, Western blot: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	The optimal dilution of the Cytokeratin 18 antibody for each application should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.