

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Moesin Antibody, Clone: [SPM562], Mouse, Monoclonal NSJ-V9046-20UG

Artikelname	Moesin Antibody, Clone: [SPM562], Mouse, Monoclonal
Artikelnummer	NSJ-V9046-20UG
Hersteller Artikelnummer	V9046-20UG
Alternativnummer	NSJ-V9046-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IF, IHC-P, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human protein was used as the immunogen for this Moesin antibody.
Produktbeschreibung	Recognizes 78kDa moesin protein. Moesin, a member of the Talin-4.1 superfamily, is a linking protein of the submembraneous actin cytoskeleton. It is expressed in variable amounts in cells of different phenotypes such as macrophages, lymphocytes, fibr...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM562]
UniProt	P26038
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT, Western blot: 1-2ug/ml, Immunofluorescence: 1-2ug/ml
Anwendungsbeschreibung	The optimal dilution of the Moesin antibody for each application should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.