

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

EMA Antibody / MUC1 / Mucin-1, Clone: [SPM132], Mouse, Monoclonal NSJ-V9047SAF-100UG

Artikelname	EMA Antibody / MUC1 / Mucin-1, Clone: [SPM132], Mouse, Monoclonal
Artikelnummer	NSJ-V9047SAF-100UG
Hersteller Artikelnummer	V9047SAF-100UG
Alternativnummer	NSJ-V9047SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Human milk fat globule membranes were used as the immunogen for this EMA antibody. This mAb reacts with the DTRP epitope in the tandem repeats.
Produktbeschreibung	Epithelial Marker Antigen is a 265-400kDa glycoprotein. The alpha subunit has cell adhesive properties. It can act both as an adhesion and an anti-adhesion protein. EMA may provide a protective layer on epithelial cells against bacterial and enzyme a...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM132]
UniProt	P15941
Reinheit	Protein G affinity chromatography

Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	The optimal dilution of the EMA antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.