

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MUC2 Antibody, Clone: [SPM296], Mouse, Monoclonal NSJ-V9049-100UG

| | |
|--------------------------|---|
| Artikelname | MUC2 Antibody, Clone: [SPM296], Mouse, Monoclonal |
| Artikelnummer | NSJ-V9049-100UG |
| Hersteller Artikelnummer | V9049-100UG |
| Alternativnummer | NSJ-V9049-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | Amino acids KYPTTTPISTTTMTPTPTGTQTPTTT from the human protein were used as the immunogen for this MUC2 antibody. |
| Produktbeschreibung | Recognizes a single glycoprotein of 520kDa, identified as mucin 2 (MUC2). This mAb shows no cross-reaction with human milk fat globule membranes, MUC1, or MUC3. Mucins are high molecular weight glycoproteins, which constitute the major component of t... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [SPM296] |
| UniProt | Q02817 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|---|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Anwendungsbeschreibung | <p>The optimal dilution of the MUC2 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9, for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p> |