

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mucin-2 Antibody, Clone: [SPM513], Mouse, Monoclonal NSJ-V9050SAF-100UG

Artikelname	Mucin-2 Antibody, Clone: [SPM513], Mouse, Monoclonal
Artikelnummer	NSJ-V9050SAF-100UG
Hersteller Artikelnummer	V9050SAF-100UG
Alternativnummer	NSJ-V9050SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Amino acids KYPTTTPISTTTMVTPTPTGTQTPTTT from the human protein were used as the immunogen for this Mucin-2 antibody.
Produktbeschreibung	Recognizes a single glycoprotein of 520kDa, identified as mucin 2 (MUC2). This mAb shows no cross-reaction with human milk fat globule membranes, MUC1, or MUC3. Its epitope has been defined as GTQTP (GlyThrGlnThrPro). Mucins are high molecular weight...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM513]
UniProt	Q02817
Reinheit	Protein G affinity chromatography

Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	<p>The optimal dilution of the Mucin-2 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p>