

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Progesterone Receptor Antibody, Clone: [SPM566], Mouse, Monoclonal NSJ-V9066IHC-7ML

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Anti-Progesterone Receptor Antibody, Clone: [SPM566], Mouse, Monoclonal                                                                                                                                                                                       |
| Artikelnummer            | NSJ-V9066IHC-7ML                                                                                                                                                                                                                                              |
| Hersteller Artikelnummer | V9066IHC-7ML                                                                                                                                                                                                                                                  |
| Alternativnummer         | NSJ-V9066IHC-7ML                                                                                                                                                                                                                                              |
| Hersteller               | NSJ Bioreagents                                                                                                                                                                                                                                               |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | IHC-P                                                                                                                                                                                                                                                         |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Recombinant human protein was used as the immunogen for this anti-Progesterone Receptor antibody.                                                                                                                                                             |
| Produktbeschreibung      | This mAb is specific to progesterone receptor and shows minimal cross-reaction with other members of the family. Progesterone receptor is expressed as two major isoforms, PR-A (81kDa) and PR-B (116kDa). Expression of PgR has been suggested to reflect... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Klon-Bezeichnung         | [SPM566]                                                                                                                                                                                                                                                      |
| UniProt                  | <a href="#">P06401</a>                                                                                                                                                                                                                                        |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Reinheit               | Protein G affinity chromatography                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Formulierung           | Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Antibody Type          | Primary Antibody                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Application Verdünnung | The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.                                                                                                                                                                                                                                                                                              |
| Anwendungsbeschreibung | The optimal dilution of the anti-Progesterone Receptor antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |