

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Bcl-2 Antibody, Clone: [SPM530], Mouse, Monoclonal NSJ-V9074IHC-7ML

Artikelname	Anti-Bcl-2 Antibody, Clone: [SPM530], Mouse, Monoclonal
Artikelnummer	NSJ-V9074IHC-7ML
Hersteller Artikelnummer	V9074IHC-7ML
Alternativnummer	NSJ-V9074IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Amino acids 41-54 (GAAPAPGIFSSQPG) from the human protein were used as the immunogen for this anti-Bcl-2 antibody.
Produktbeschreibung	This antibody recognizes a protein of 25-26kDa, identified as the Bcl-2 alpha oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of bcl-2 alpha oncoprotein inhibits the programmed cell death (apoptosis). In most follicular ...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM530]
UniProt	P10415
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	The optimal dilution of the anti-Bcl-2 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.