

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Bcl-2 Antibody, Clone: [SPM530], Mouse, Monoclonal NSJ-V9074SAF-100UG

Artikelname	Anti-Bcl-2 Antibody, Clone: [SPM530], Mouse, Monoclonal
Artikelnummer	NSJ-V9074SAF-100UG
Hersteller Artikelnummer	V9074SAF-100UG
Alternativnummer	NSJ-V9074SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	FACS, IF, IHC-P, WB
Spezies Reaktivität	Human
Immunogen	Amino acids 41-54 (GAAPAPGIFSSQPG) from the human protein were used as the immunogen for this anti-Bcl-2 antibody.
Produktbeschreibung	This antibody recognizes a protein of 25-26kDa, identified as the Bcl-2 alpha oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of bcl-2 alpha oncoprotein inhibits the programmed cell death (apoptosis). In most follicular ...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM530]
UniProt	P10415
Reinheit	Protein G affinity chromatography

Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Verdünnung	Flow cytometry: 1-2ug/10 ⁶ cells, Immunofluorescence: 1-2ug/ml, Western blot: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	The optimal dilution of the anti-Bcl-2 antibody for each application should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.