

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Bcl-X Antibody, Clone: [SPM165], Mouse, Monoclonal NSJ-V9075-100UG

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Anti-Bcl-X Antibody, Clone: [SPM165], Mouse, Monoclonal                                                                                                                                                                                                       |
| Artikelnummer            | NSJ-V9075-100UG                                                                                                                                                                                                                                               |
| Hersteller Artikelnummer | V9075-100UG                                                                                                                                                                                                                                                   |
| Alternativnummer         | NSJ-V9075-100UG                                                                                                                                                                                                                                               |
| Hersteller               | NSJ Bioreagents                                                                                                                                                                                                                                               |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | IHC-P                                                                                                                                                                                                                                                         |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Amino acids 3-14 (QSNRELVVDFLS) from the human protein were used as the immunogen for this anti-Bcl-X antibody.                                                                                                                                               |
| Produktbeschreibung      | Recognizes a protein of 27kDa, identified as the Bcl-X protein. This mAb shows no cross-reaction with Bcl-2 or Bax protein. Bcl-X has two isoforms, Bcl-XL (long), a 241 amino acid protein which suppresses cell death. And Bcl-XS (short), a 178 amino a... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Klon-Bezeichnung         | [SPM165]                                                                                                                                                                                                                                                      |
| UniProt                  | <a href="#">Q07817</a>                                                                                                                                                                                                                                        |
| Reinheit                 | Protein G affinity chromatography                                                                                                                                                                                                                             |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Formulierung           | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Antibody Type          | Primary Antibody                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Anwendungsbeschreibung | <p>The optimal dilution of the anti-Bcl-X antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p> |