

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

gp100 Antibody / PMEL17, Clone: [SPM142], Mouse, Monoclonal NSJ-V9078SAF-100UG

Artikelname	gp100 Antibody / PMEL17, Clone: [SPM142], Mouse, Monoclonal
Artikelnummer	NSJ-V9078SAF-100UG
Hersteller Artikelnummer	V9078SAF-100UG
Alternativnummer	NSJ-V9078SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P, WB
Spezies Reaktivität	Human
Immunogen	Extract of pigmented melanoma metastases from lymph nodes was used as the immunogen for this gp100 antibody.
Produktbeschreibung	By immunohistochemistry, it specifically recognizes a protein in melanocytes and melanomas. This mAb reacts with junctional and blue nevus cells and variably with fetal and neonatal melanocytes. Intradermal nevi, normal adult melanocytes, and non-mel...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM142]
UniProt	P40967
Reinheit	Protein G affinity chromatography

Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Verdünnung	Western blot: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	The optimal dilution of the gp100 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.