

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-TG Antibody, Clone: [SPM517], Mouse, Monoclonal NSJ-V9084-100UG

Artikelname	Anti-TG Antibody, Clone: [SPM517], Mouse, Monoclonal
Artikelnummer	NSJ-V9084-100UG
Hersteller Artikelnummer	V9084-100UG
Alternativnummer	NSJ-V9084-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Human thyroid follicular cells were used as the immunogen for this anti-TG antibody.
Produktbeschreibung	Thyroglobulin (TG) is a 660kDa dimeric pre-protein with multiple glycosylation sites. It is produced by and processed within the thyroid gland to produce the hormone thyroxine and triiodothyronine. Prior to forming dimers, thyroglobulin monomers under...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM517]
UniProt	<a href="#">P01266</a>
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	<p>The optimal dilution of the anti-TG antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p>