

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-TG Antibody, Clone: [SPM517], Mouse, Monoclonal NSJ-V9084SAF-100UG

| | |
|--------------------------|--|
| Artikelname | Anti-TG Antibody, Clone: [SPM517], Mouse, Monoclonal |
| Artikelnummer | NSJ-V9084SAF-100UG |
| Hersteller Artikelnummer | V9084SAF-100UG |
| Alternativnummer | NSJ-V9084SAF-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | Human thyroid follicular cells were used as the immunogen for this anti-TG antibody. |
| Produktbeschreibung | Thyroglobulin (TG) is a 660kDa dimeric pre-protein with multiple glycosylation sites. It is produced by and processed within the thyroid gland to produce the hormone thyroxine and triiodothyronine. Prior to forming dimers, thyroglobulin monomers under... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [SPM517] |
| UniProt | P01266 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|--|
| Formulierung | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Anwendungsbeschreibung | <p>The optimal dilution of the anti-TG antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p> |