

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-TGF-alpha Antibody, Clone: [SPM542], Mouse, Monoclonal NSJ-V9086-20UG

| | |
|--------------------------|---|
| Artikelname | Anti-TGF-alpha Antibody, Clone: [SPM542], Mouse, Monoclonal |
| Artikelnummer | NSJ-V9086-20UG |
| Hersteller Artikelnummer | V9086-20UG |
| Alternativnummer | NSJ-V9086-20UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | Amino acids 34-43 (PPVAAAVVSH) from the human protein were used as the immunogen for this anti-TGF alpha antibody. |
| Produktbeschreibung | This antibody reacts with the TGF alpha and shows no cross-reaction with EGF and the neuropeptide synenkephalin. TGFa is a growth factor with 33% homology to EGF, binds to EGFR, activates tyrosine phosphorylation of the receptor, and stimulates cell ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [SPM542] |
| UniProt | P01135 |
| Reinheit | Precipitated followed by dialysis in 1X PBS |

| | |
|------------------------|--|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Anwendungsbeschreibung | The optimal dilution of the anti-TGF-alpha antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes. |