

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-PGP9.5 Antibody, Clone: [SPM574], Mouse, Monoclonal NSJ-V9096-100UG

Artikelname	Anti-PGP9.5 Antibody, Clone: [SPM574], Mouse, Monoclonal
Artikelnummer	NSJ-V9096-100UG
Hersteller Artikelnummer	V9096-100UG
Alternativnummer	NSJ-V9096-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IF, IHC-P, WB
Spezies Reaktivität	Human, Rat
Immunogen	Native UchL1/PGP9.5 protein isolated from human brain was used as the immunogen for this anti-PGP9.5 antibody.
Produktbeschreibung	This mAb reacts with a protein of 20-30kDa, identified as PGP9.5, also known as ubiquitin carboxyl-terminal hydrolase-1 (UchL1). Initially, PGP9.5 expression in normal tissues was reported in neurons and neuroendocrine cells but later it was found in...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM574]
UniProt	P09936
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Western blot: 1-2ug/ml,Immunofluorescence: 1-2ug/ml,Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	The optimal dilution of the anti-PGP9.5 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.