

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Carbonic Anhydrase IX Antibody, Clone: [SPM487], Mouse, Monoclonal NSJ-V9101-100UG

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Carbonic Anhydrase IX Antibody, Clone: [SPM487], Mouse, Monoclonal                                                                                                                                                                                            |
| Artikelnummer            | NSJ-V9101-100UG                                                                                                                                                                                                                                               |
| Hersteller Artikelnummer | V9101-100UG                                                                                                                                                                                                                                                   |
| Alternativnummer         | NSJ-V9101-100UG                                                                                                                                                                                                                                               |
| Hersteller               | NSJ Bioreagents                                                                                                                                                                                                                                               |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | IHC-P                                                                                                                                                                                                                                                         |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | The microsomal fraction of human renal cortical tissue homogenate was used as the immunogen for this Carbonic Anhydrase IX antibody.                                                                                                                          |
| Produktbeschreibung      | Recognizes a glycoprotein of ~200kDa, identified as carbonic anhydrase IX (CAIX/CA9/gp200). Its epitope resides in the carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate determinants, such as the Lewis bloo... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Klon-Bezeichnung         | [SPM487]                                                                                                                                                                                                                                                      |
| UniProt                  | <a href="#">Q16790</a>                                                                                                                                                                                                                                        |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
|------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Reinheit               | Protein G affinity chromatography                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Formulierung           | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Antibody Type          | Primary Antibody                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Anwendungsbeschreibung | <p>The optimal dilution of the Carbonic Anhydrase IX antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p> |