

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-CD44 Antibody, Clone: [SPM544], Mouse, Monoclonal NSJ-V9105-100UG

Artikelname	Anti-CD44 Antibody, Clone: [SPM544], Mouse, Monoclonal
Artikelnummer	NSJ-V9105-100UG
Hersteller Artikelnummer	V9105-100UG
Alternativnummer	NSJ-V9105-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P, WB
Spezies Reaktivität	Human
Immunogen	Stimulated human leukocytes were used as the immunogen for this anti-CD44 antibody.
Produktbeschreibung	Recognizes a cell surface glycoprotein of 80-95kDa (CD44) on lymphocytes, monocytes, and granulocytes. Its epitope is resistant to digestion by trypsin and chymotrypsin. The CD44 family of glycoproteins exists in a number of variant isoforms, the mos...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM544]
UniProt	P16070
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Western blot: 1-2ug/ml,Immunohistochemistry (FFPE): 0.25-0.5ug/ml for 30 min at RT
Anwendungsbeschreibung	The optimal dilution of the anti-CD44 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.