

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Hepatocyte Antibody, Clone: [SPM582], Mouse, Monoclonal NSJ-V9119-20UG

Artikelname	Hepatocyte Antibody, Clone: [SPM582], Mouse, Monoclonal
Artikelnummer	NSJ-V9119-20UG
Hersteller Artikelnummer	V9119-20UG
Alternativnummer	NSJ-V9119-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Extract of a formalin-fixed rejected-allograft of human liver was used as the immunogen for this Hepatocyte antibody.
Produktbeschreibung	Hepatocyte Specific Antigen, also called Hepatocyte Paraffin 1 or Hep Par 1, localizes to the mitochondria of hepatocytes. It is a sensitive marker for distinguishing hepatocellular carcinomas (HCC) from other metastatic carcinomas as well as cholang...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM582]
UniProt	Not Known
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	The optimal dilution of the Hepatocyte antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.