

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Recombinant Pan-CK Antibody Cocktail, Clone: [Cocktail rPCK/6750], Mouse, Monoclonal NSJ-V9362-100UG**

|                          |   |
|--------------------------|---|
| Artikelname              | Recombinant Pan-CK Antibody Cocktail, Clone: [Cocktail rPCK/6750],<br>Mouse, Monoclonal   |
| Artikelnummer            | NSJ-V9362-100UG   |
| Hersteller Artikelnummer | V9362-100UG   |
| Alternativnummer         | NSJ-V9362-100UG   |
| Hersteller               | NSJ Bioreagents   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FACS, IF, IHC-P   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Crude cytokeratin extract prepared from RT-4 and MCF-7 cells was used as the immunogen for the recombinant Pan-CK antibody.   |
| Produktbeschreibung      | Anti-cytokeratin clone rPCK/6750 demonstrates a broad spectrum of cytokeratin reactivity. In normal tissues, rPCK/6750 is reactive with most epithelial types, including bile ducts and hepatocytes in liver, bladder epithelium, breast ducts, bronchial ... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [Cocktail rPCK/6750]  |
| Reinheit                 | Protein A/G affinity  |

|                        |  |
|------------------------|--|
| Formulierung           | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide                                  |
| Antibody Type          | Primary Antibody   |
| Application Verdünnung | Flow cytometry: 1-2ug/million cells, Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml |
| Anwendungsbeschreibung | Optimal dilution of the recombinant Pan-CK antibody should be determined by the researcher.              |