

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Recombinant Human LR3-IGF-1 EBT-EPT077

Artikelname	Recombinant Human LR3-IGF-1
Artikelnummer	EBT-EPT077
Hersteller Artikelnummer	EPT077
Alternativnummer	EBT-EPT077-50
Hersteller	ELK Biotechnology
Kategorie	Proteine/Peptide
Produktbeschreibung	Recombinant Human LR3 Insulin-Like Growth Factor-I is produced by our E.coli expression system and the target gene encoding Gly49-Ala118 is expressed....
Molekulargewicht	Molecular weight: 9.1 KDa. Apparent molecular weight: 11 KDa, reducing conditions
UniProt	<a href="#">P05019</a>
Reinheit	Greater than 95% as determined by reducing SDS-PAGE.

Anwendungsbeschreibung

Redissolve: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in 50mM Acetic Acid. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.. Endotoxin: Less than 0.05 ng/µg (0.5 EU/µg) as determined by LAL test. Background: Insulin-like growth factor I (IGF1) belongs to the family of insulin-like growth factors that are structurally homologous to proinsulin. Mature IGFs are generated by proteolytic processing of inactive precursor proteins, which contains the N- and C-terminal propeptide regions. Mature human IGF-I consisting of 70 amino acids has 94% identity with mouse IGF-I and exhibits cross-species activity. IGF-1 binds IGF-IR, IGF-IIR, and the insulin receptor and plays a key role in cell cycle progression, cell proliferation and tumor progression. IGF-1 expression is regulated by growth hormone. R3 IGF-1 is an 83 amino acid analog of IGF-1 comprising the complete human IGF-1 sequence with the substitution of an Arg (R) for the Glu(E) at position three, hence R3, and a 13 amino acid extension peptide at the N terminus. R3 IGF-1 has been produced with the purpose of increasing biological activity. R3 IGF-1 is significantly more potent than human IGF-I in vitro