

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant Mouse IL-12 EBT-EPT228

Artikelname	Recombinant Mouse IL-12
Artikelnummer	EBT-EPT228
Hersteller Artikelnummer	EPT228
Alternativnummer	EBT-EPT228-10
Hersteller	ELK Biotechnology
Kategorie	Proteine/Peptide
Produktbeschreibung	Recombinant Mouse Interleukin-12 is produced by our Mammalian expression system and the target gene encoding Met23-Ser335&Arg23-Ala215 is expressed....
Molekulargewicht	Molecular weight: 35.8&21.7 KDa. Apparent molecular weight: 40-55&20-28 KDa, reducing conditions
UniProt	P43432
Reinheit	Greater than 90% as determined by reducing SDS-PAGE.

Anwendungsbeschreibung	Redissolve: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Endotoxin: Less than 0.001 ng/µg (0.01 EU/µg) as determined by LAL test. Background: Mouse IL-12 is a heterodimeric pleiotropic cytokine made up of a 40 kDa (p40) subunit and a 35 kDa (p35) subunit. Human and mouse IL-12 share 70% and 60% amino acid sequence identity in their p40 and p35 subunits, respectively. While mouse IL-12 is active on both human and mouse cells, human IL-12 is not active on murine cells. It is involved in the differentiation of naive T cells into Th1 cells. It is known as a T cell-stimulating factor, which can stimulate the growth and function of T cells. It stimulates the production of interferon-gamma (IFN-gamma) and tumor necrosis factor-alpha (TNF-alpha) from T cells and natural killer (NK) cells, and reduces IL-4 mediated suppression of IFN-gamma. T cells that produce IL-12 have a coreceptor, CD30, which is associated with IL-12 activity. IL-12 plays an important role in the activities of natural killer cells and T lymphocytes. IL-12 mediates enhancement of the cytotoxic activity of NK cells and CD8+ cytotoxic T lymphocytes.
------------------------	--