

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Recombinant Rabbit Monoclonal Antibody to Chromogranin A / CHGA (Neuroendocrine Marker)(Clone : CHGA/1815R), Clone: [CHGA/1815R] ABI-12-1018-20**

|                          |   |
|--------------------------|---|
| Artikelname              | Recombinant Rabbit Monoclonal Antibody to Chromogranin A / CHGA (Neuroendocrine Marker)(Clone : CHGA/1815R), Clone: [CHGA/1815R]  |
| Artikelnummer            | ABI-12-1018-20  |
| Hersteller Artikelnummer | 12-1018-20  |
| Alternativnummer         | ABI-12-1018-20-20UG   |
| Hersteller               | Abeomics  |
| Kategorie                | Sonstiges   |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant full-length human CHGA protein  |
| Produktbeschreibung      | Chromogranin A is present in neuroendocrine cells throughout the body, including the neuroendocrine cells of the large and small intestine, adrenal medulla and pancreatic islets. It is an excellent marker for carcinoid tumors, pheochromocytomas, para... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [CHGA/1815R]  |
| NCBI                     | <a href="#">1113</a>  |
| UniProt                  | <a href="#">P10645</a>  |

|                        |  |
|------------------------|--|
| Reinheit               | Purified Ab with BSA and Azide at 200ug/ml   |
| Formulierung           | Purified   |
| Target-Kategorie       | Chromogranin A / CHGA (Neuroendocrine Marker)  |
| Antibody Type          | Recombinant Antibody   |
| Application Verdünnung | Immunohistochemistry (Formalin-fixed) (1-2µg/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes), |