

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Tyrosinase (Melanoma Marker) Recombinant Rabbit Monoclonal Antibody (Clone:TYR/2024R), Clone: [TYR/2024R] ABI-12-1262

|                          |   |
|--------------------------|---|
| Artikelname              | Anti-Tyrosinase (Melanoma Marker) Recombinant Rabbit Monoclonal Antibody (Clone:TYR/2024R), Clone: [TYR/2024R]  |
| Artikelnummer            | ABI-12-1262   |
| Hersteller Artikelnummer | 12-1262   |
| Alternativnummer         | ABI-12-1262-100UG   |
| Hersteller               | Abeomics  |
| Wirt                     | Rabbit  |
| Kategorie                | Sonstiges   |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant tyrosinase protein  |
| Produktbeschreibung      | Recognizes a cluster of proteins between 70-80kDa, identified as tyrosinase. Occasionally a minor band at 55kDa is also detected. This MAb shows no cross-reaction with MAGE-1 and tyrosinase-related protein 1, TRP-1/gp75. Tyrosinase is a copper-contai... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [TYR/2024R]   |
| NCBI                     | <a href="#">7299</a>  |
| UniProt                  | <a href="#">P14679</a>  |

|                        |  |
|------------------------|--|
| Reinheit               | Protein A/G  |
| Formulierung           | Purified   |
| Antibody Type          | Recombinant Antibody   |
| Application Verdünnung | Immunohistochemistry (Formalin-fixed) (1-2µg/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes), |