

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Folate Binding Protein Antibody, Unconjugated, Goat, Polyclonal Preis auf Anfrage BYT-ORB344183

| | |
|--------------------------|---|
| Artikelname | Folate Binding Protein Antibody, Unconjugated, Goat, Polyclonal Preis auf Anfrage |
| Artikelnummer | BYT-ORB344183 |
| Hersteller Artikelnummer | orb344183 |
| Alternativnummer | BYT-ORB344183-25 |
| Hersteller | Biorbyt |
| Wirt | Goat |
| Kategorie | Antikörper |
| Applikation | ELISA, IP, WB |
| Spezies Reaktivität | Bovine |
| Immunogen | Folate Binding Protein [Bovine Milk] |
| Konjugation | Unconjugated |
| Produktbeschreibung | Folate Binding Protein antibody... |
| Klonalität | Polyclonal |
| Konzentration | 1 mg/ml |
| NCBI | 001193456 |
| UniProt | P02702 |

| | |
|------------------------|--|
| Puffer | Preservative: 0.01% (w/v) Sodium Azide. Stabilizer: None, Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Reinheit | Anti-FOLATE BINDING PROTEIN is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum as well as purified and partially purified Folate Binding Protein [Bovine Milk]. Cross reactivity against Folate Binding Protein from other tissues and species may occur but have not been specifically determined. |
| Formulierung | Liquid (sterile filtered) |
| Application Verdünnung | ELISA: 1:5,000 - 1:20,000, IP: 1:100, WB: 1:500 - 1:5,000 |
| Anwendungsbeschreibung | Application Notes: FOLATE BINDING PROTEIN Antibody has been assayed against 1.0 ug of Folate Binding Protein [Bovine Milk] in a standard ELISA using Peroxidase conjugated Affinity Purified anti-Goat IgG [H&L] (Rabbit) and (ABTS (2,2-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:10,000 to 1:40,000 of the reconstitution concentration is suggested for this product. Highest levels of antigen binding and antibody reactivity were observed using NuncO brand MaxiSorp immunoplates |