

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Acetylcholinesterase Antibody Peroxidase Conjugated, HRP, Goat, Polyclonal Preis auf Anfrage BYT-ORB344266

Artikelname	Acetylcholinesterase Antibody Peroxidase Conjugated, HRP, Goat, Polyclonal Preis auf Anfrage
Artikelnummer	BYT-ORB344266
Hersteller Artikelnummer	orb344266
Alternativnummer	BYT-ORB344266-25
Hersteller	Biorbyt
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA, IP, WB
Spezies Reaktivität	Bovine
Immunogen	Acetylcholinesterase [Bovine Erythrocytes]
Konjugation	HRP
Produktbeschreibung	Acetylcholinesterase antibody (Peroxidase)...
Klonalität	Polyclonal
Konzentration	1 mg/mL
NCBI	001069688
UniProt	P23795

Puffer	Preservative: 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!. Stabilizer: 10 mg/mL Bovine Serum Albumin (rAlbumin) - Immunoglobulin and Protease free, Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	Anti-Acetylcholinesterase (Bovine Erythrocytes) antibody is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum as well as purified and partially purified Acetylcholinesterase [Bovine Erythrocytes]. Cross reactivity against Acetylcholinesterase from other sources is unknown.
Formulierung	Liquid (sterile filtered)
Application Verdünnung	ELISA: 1:2,000 - 1:10,000, IP: 1:100, WB: 1:500 - 1:2,500
Anwendungsbeschreibung	Application Notes: Anti-Acetylcholinesterase (Bovine Erythrocytes) antibody has been tested by western blot and is suitable to be assayed against 1.0 µg of Acetylcholinesterase [Bovine Erythrocytes] in a standard capture ELISA using ABTS (2,2-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:1,000 of the reconstitution concentration is suggested for this product