

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

BUBR1 Kinase Antibody, IgG1, Clone: [8G1], Unconjugated, Mouse, Monoclonal Preis auf Anfrage BYT-ORB344429

Artikelname	BUBR1 Kinase Antibody, IgG1, Clone: [8G1], Unconjugated, Mouse, Monoclonal Preis auf Anfrage
Artikelnummer	BYT-ORB344429
Hersteller Artikelnummer	orb344429
Alternativnummer	BYT-ORB344429-100
Hersteller	Biorbyt
Wirt	Mouse
Kategorie	Antikörper
Applikation	ELISA, IF, IP, WB
Spezies Reaktivität	Human
Immunogen	This protein A purified monoclonal antibody was produced by repeated immunizations with a recombinant protein corresponding to amino acid residues 1-350 of human BUBR1 Kinase protein.
Konjugation	Unconjugated
Produktbeschreibung	BUBR1 Kinase Kinase antibody...
Klonalität	Monoclonal
Konzentration	1.0 mg/mL
Klon-Bezeichnung	[8G1]

Isotyp	IgG1
NCBI	001202
UniProt	O60566
Puffer	Preservative: 0.01% (w/v) Sodium Azide. Stabilizer: None, Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This Protein A purified antibody is directed against human BUBR1 Kinase protein. The product was purified from tissue culture supernatant by chromatography. This antibody has only been tested on human cells. Reactivity against homologues from other sources is not known.
Formulierung	Liquid (sterile filtered)
Application Verdünnung	ELISA: 1:5,000 - 1:20,000, IF: 1:200 - 1:1,000, IP: 1:100, WB: 1:200 - 1:1,000
Anwendungsbeschreibung	Application Notes: This protein A purified antibody has been tested for use in immunoprecipitation, immunofluorescence staining and western blot and is capable of detecting endogenous protein. Specific conditions for reactivity should be optimized by the end user. Expect a predominant band at ~ 120 kDa corresponding to full-length protein by western blotting in the appropriate cell lysate or extract. Higher MW bands may be seen that may be due to hyper-phosphorylation of the protein. The use of HeLa whole cell lysates is recommended as a positive control. For IF microscopy use cells grown on cover slips fixed with 3.5% paraformaldehyde in PBS at pH 6.8. Permeabilize fixed cells with 0.2% Triton X-100 in 25 mM Tris Cl, pH 7.4 containing 0.1% BSA