

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

LGR4 Antibody, IgG2b, Clone: [6G8.B3.G5.C3], Unconjugated, Mouse, Monoclonal Preis auf Anfrage BYT-ORB344461

Artikelname	LGR4 Antibody, IgG2b, Clone: [6G8.B3.G5.C3], Unconjugated, Mouse, Monoclonal Preis auf Anfrage
Artikelnummer	BYT-ORB344461
Hersteller Artikelnummer	orb344461
Alternativnummer	BYT-ORB344461-1
Hersteller	Biorbyt
Wirt	Mouse
Kategorie	Antikörper
Applikation	ELISA, IHC, WB
Spezies Reaktivität	Human, Monkey, Primate
Immunogen	This LGR4 antibody was produced by repeated immunizations with a synthetic peptide corresponding to an internal region of human LGR4 protein. The hybridoma was produced by fusing BALB/c mouse splenocytes and mouse myeloma SP2/O cells using conventional technology.
Konjugation	Unconjugated
Produktbeschreibung	LGR4 antibody...
Klonalität	Monoclonal
Konzentration	1 mg/ml

Klon-Bezeichnung	[6G8.B3.G5.C3]
Isotyp	IgG2b
NCBI	157694513
UniProt	Q8N537
Puffer	Preservative: 0.01% (w/v) Sodium Azide. Stabilizer: None, Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	Anti-LGR4 Antibody product was purified from concentrated tissue culture supernate by Protein A chromatography. This antibody is specific for human LGR4 protein. A BLAST analysis was used to suggest cross-reactivity with LGR4 from chimpanzee, orangutan and macaque based on 100% homology with the immunizing sequence. Cross-reactivity with LGR4 from other sources has not been determined.
Formulierung	Liquid (sterile filtered)
Application Verdünnung	ELISA: 1:20,000 - 1:100,000, IHC: 5 ug/mL, WB: 1:500 - 1:3,000
Anwendungsbeschreibung	Application Notes: Anti-LGR4 antibody has been tested by ELISA and western blotting, and is suitable in immunohistochemistry. Expect a band approximately 102 kDa in size corresponding to LGR4 protein by western blotting in the appropriate cell lysate or extract. Specific conditions for reactivity should be optimized by the end user. Use formalin-fixed paraffin-embedded sections for immunohistochemistry. No pre-treatment of sample is required