

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Nectin-3, Mouse, mAb 103-A1 Antibody, Unconjugated, Monoclonal BYT-ORB1109146

| | |
|--------------------------|--|
| Artikelname | Nectin-3, Mouse, mAb 103-A1 Antibody, Unconjugated, Monoclonal |
| Artikelnummer | BYT-ORB1109146 |
| Hersteller Artikelnummer | orb1109146 |
| Alternativnummer | BYT-ORB1109146-100 |
| Hersteller | Biorbyt |
| Kategorie | Antikörper |
| Applikation | FC, Functional Studies, IA, IF, IHC-Fr, IP |
| Spezies Reaktivität | Mouse |
| Immunogen | Recombinant mouse nectin-3 extracellular domain (50-400 aa) fused to IgG Fc |
| Konjugation | Unconjugated |
| Produktbeschreibung | Mouse monoclonal antibody to Nectin-3... |
| Klonalität | Monoclonal |
| Puffer | 0.2 ml (100 µg/ml) 0.2 µm filtered antibody solution in PBS, containing 0.1% bovine serum albumin and 0.02% sodium azide |
| Formulierung | 1 ml (100 µg/ml) 0.2 µm filtered antibody solution in PBS, containing 0.1% bovine serum albumin and 0.02% sodium azide |

| | |
|------------------------|---|
| Application Verdünnung | For immunohistochemistry and flow cytometry, dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50. |
| Anwendungsbeschreibung | Application Notes: IF: fixed in 2% paraformaldehyde in PBS optionally followed by methanol/acetone fixation (Ref 2-7) FC: 1 ug per 1 x 10 ⁶ cells FS: injection of 1ug into lumen of seminiferous tubules (Ref 7) IA: Indirect ELISA on 0.5µg/ml mouse nectin-3 coated per well. Positive signal with 2µg/ml antibody. IF: Cells are fixed in 2 % paraformaldehyde in PBS, optionally followed by methanol/acetone fixation (Ref 1-6) IP: 1-2 ug per 100-500 ug of total protein (1 ml of cell lysate) |