

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### ODC1 Antibody, Clone: [ODC1/485], Unconjugated, Mouse, Monoclonal BYT-ORB1252723

|                          |                                                                              |
|--------------------------|------------------------------------------------------------------------------|
| Artikelname              | ODC1 Antibody, Clone: [ODC1/485], Unconjugated, Mouse, Monoclonal            |
| Artikelnummer            | BYT-ORB1252723                                                               |
| Hersteller Artikelnummer | orb1252723                                                                   |
| Alternativnummer         | BYT-ORB1252723-100                                                           |
| Hersteller               | Biorbyt                                                                      |
| Wirt                     | Mouse                                                                        |
| Kategorie                | Antikörper                                                                   |
| Applikation              | FC, IF, IHC, WB                                                              |
| Spezies Reaktivität      | Human, Rat                                                                   |
| Immunogen                | Recombinant human ODC-1 protein was used as the immunogen for this antibody. |
| Konjugation              | Unconjugated                                                                 |
| Produktbeschreibung      | ODC1 Antibody...                                                             |
| Klonalität               | Monoclonal                                                                   |
| Konzentration            | 0.2 mg/mL                                                                    |
| Klon-Bezeichnung         | [ODC1/485]                                                                   |
| UniProt                  | <a href="#">P11926</a>                                                       |
| Puffer                   | PBS with 0.1 mg/ml rAlbumin and 0.05% sodium azide                           |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Formulierung           | Liquid                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| Target-Kategorie       | ODC1                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Anwendungsbeschreibung | <p>Application Notes: Flow Cytometry: 0.5-1 ug/million cellsIF: 0.5-1 µg/mLWB: 0.5-1 µg/mL IHC (FFPE): 0.5-1 µg/mL for 30 minutes at RT (1)Prediluted format : incubate for 30 min at RT (2)The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the antibody to be titrated up or down for optimal performance.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min</p> |