**BIOZOL**

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

**E-Mail:** info@biozol.de

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

# Product Datasheet

## Rabbit IgG anti-Llama IgG (VHH)-unconj., MinX none

## DNA-SEC-182476

| | |
|---|---|
| Artikelname | Rabbit IgG anti-Llama IgG (VHH)-unconj., MinX none |
| Artikelnummer | DNA-SEC-182476 |
| Hersteller Artikelnummer | SEC-182476 |
| Alternativnummer | DNA-SEC-182476 |
| Hersteller | dianova |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | ELISA,WB |
| Spezies Reaktivität | Llama |
| Immunogen | This antibody was prepared from whole rabbit serum produced by repeated immunizations with a VHH camelid domain protein. |
| Konjugation | Unconjugated |
| Format | IgG |
| Spezifität | IgG (VHH) |
| Minimale Kreuzreaktivität (MinX) | no cross-adsorbtion |

| | |
|---|---|
| Produktbeschreibung | Rabbit Anti-VhH antibody can be used to detect single domain VhH antibodies. The single-domain antibody (sdAb) is a small (12 - 14 kDa) antibody fragment that consists of a monomeric variable domain derived from the heavy chain, also called a VhH ant... |
| Klonalität | Polyclonal |
| Konzentration | 1.3 mg/mL |
| Isotyp | Ig |
| Puffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Reinheit | Anti-VHH is directed against the llama VHH protein fragment. The product was purified from monospecific antiserum by protein A chromatography. Anti-VHH antibody detects recombinant VHH proteins, native Llama IgG2 and native Llama IgG3. |
| Formulierung | Liquid (sterile filtered) |
| Formel | 20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3 |
| Target-Kategorie | Llama |
| Antibody Type | Secondary Antibody |
| Application Verdünnung | ELISA Dilution: 1:10,000-1:50,000, Western Blot Dilution: 1:1,000-1:5,000 |
| Anwendungsbeschreibung | Anti-VHH antibody has been tested by ELISA and Western blot. Specific conditions for reactivity should be optimized by the end user. Some cross-reactivity to E.coli proteins may be observed. |