**BIOZOL**

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

**E-Mail:** info@biozol.de

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

# Product Datasheet

## Rabbit IgG anti-Bovine IgG (H+L)-unconj., MinX none

## DNA-SEC-182482

| | |
|---|---|
| Artikelname | Rabbit IgG anti-Bovine IgG (H+L)-unconj., MinX none |
| Artikelnummer | DNA-SEC-182482 |
| Hersteller Artikelnummer | SEC-182482 |
| Alternativnummer | DNA-SEC-182482 |
| Hersteller | dianova |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | ELISA,IHC,WB |
| Spezies Reaktivität | Bovine |
| Immunogen | Bovine IgG whole molecule |
| Konjugation | Unconjugated |
| Format | IgG |
| Spezifität | IgG (H+L) |
| Minimale Kreuzreaktivität (MinX) | no cross-adsorbtion |

| | |
|---|---|
| Produktbeschreibung | Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglu... |
| Klonalität | Polyclonal |
| Konzentration | 10.0 mg/mL |
| Isotyp | Ig |
| Puffer | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Reinheit | This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Bovine IgG and Bovine Serum. |
| Formulierung | Lyophilized |
| Formel | 10 mM NaPO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3 |
| Target-Kategorie | Bovine |
| Antibody Type | Secondary Antibody |
| Application Verdünnung | ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000 |
| Anwendungsbeschreibung | Anti-Bovine IgG antibody is suitable for ELISA, western blot, and immunohistochemistry, as well as other assays requiring lot-to-lot consistency. |