BIOZOL

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Chicken IgG anti-Goat IgG (H+L)-Alk. Phos., MinX none

DNA-SEC-182895

Artikelname	Chicken IgG anti-Goat IgG (H+L)-Alk. Phos., MinX none
Artikelnummer	DNA-SEC-182895
Hersteller Artikelnummer	SEC-182895
Alternativnummer	DNA-SEC-182895
Hersteller	dianova
Wirt	Gallus
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Goat
Immunogen	Goat IgG whole molecule
Konjugation	Alk. Phos.
Format	IgG
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion

Produktbeschreibung	Anti-Goat IgG Alkaline Phosphatase Antibody generated in chicken detects goat IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacte...
Klonalität	Polyclonal
Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Chicken Serum, Goat IgG and Goat Serum.
Formulierung	Liquid (sterile filtered)
Formel	50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN3
Target-Kategorie	Goat
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:2,000 - 1:10,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500
Anwendungsbeschreibung	This product has been assayed against 1.0 ug of Goat IgG in a standard capture ELISA using pNPP (p-nitrophenyl phosphate) code # NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:500 to 1:2,000 of the stated concentration is suggested for this product.