BIOZOL

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Human IgG+IgM+IgA (H+L)-unconj., MinX none

DNA-SEC-182989

Artikelname	Goat IgG anti-Human IgG+IgM+IgA (H+L)-unconj., MinX none
Artikelnummer	DNA-SEC-182989
Hersteller Artikelnummer	SEC-182989
Alternativnummer	DNA-SEC-182989
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Human
Immunogen	Human IgG, IgA and IgM whole molecules
Konjugation	Unconjugated
Format	IgG
Spezifität	IgG+IgM+IgA (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion

Produktbeschreibung	Anti-Human IgG IgA IgM (H&L) Antibody generated in goat detects human (heavy and light chain) immunoglobulin G, A, and M. Immunoglobulin G binds to antigens and can neutralize or opsonize targets, and are predominantly involved in the secondary immun...
Klonalität	Polyclonal
Konzentration	2.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from polyspecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Human IgG, Human IgA and Human IgM. This reagent is suitable for the detection of all Human immunoglobulin subclasses and chain combinations.
Formulierung	Liquid (sterile filtered)
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3
Target-Kategorie	Human
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:28,800 - 1:48,800, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Anwendungsbeschreibung	Anti-Human IgG IgA IgM antibody has been tested by ELISA and dot blot and is suitable for western blot and immunohistochemistry, as well as other assays requiring lot-to-lot consistency.