BIOZOL

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Mouse IgG+IgM (H+L)-HRPO, MinX none

DNA-SEC-183135

Artikelname	Goat IgG anti-Mouse IgG+IgM (H+L)-HRPO, MinX none
Artikelnummer	DNA-SEC-183135
Hersteller Artikelnummer	SEC-183135
Alternativnummer	DNA-SEC-183135
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Mouse
Immunogen	Anti-Mouse IgG and IgM whole molecule was produced by repeated immunization with Mouse IgG and IgM whole molecule in goat.
Konjugation	HRPO
Format	IgG
Spezifität	IgG+IgM (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion

Produktbeschreibung	Anti-Mouse IgG and IgM whole molecule peroxidase conjugated antibody generated in goat detects specifically Mouse IgG and IgM whole molecule. This secondary peroxidase conjugated antibody anti-mouse is ideal for investigators who routinely perform ti...
Klonalität	Polyclonal
Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from polyspecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc agains No reaction was observed against bovine, horse or human serum proteins.
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target-Kategorie	Mouse
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:40,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:2,000 - 1:10,000
Anwendungsbeschreibung	Anti-Mouse IgG and IgM Peroxidase conjugated antibody has been tested by ELISA and is suitable for highly specific immunological methods requiring extremely low background levels, lot-to-lot consistency, high titer and specificity.