BIOZOL

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Sheep IgG anti-Mouse IgG (H+L)-Alk. Phos., MinX none

DNA-SEC-183327

Artikelname	Sheep IgG anti-Mouse IgG (H+L)-Alk. Phos., MinX none
Artikelnummer	DNA-SEC-183327
Hersteller Artikelnummer	SEC-183327
Alternativnummer	DNA-SEC-183327
Hersteller	dianova
Wirt	Sheep
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Mouse
Immunogen	Mouse IgG whole molecule
Konjugation	Alk. Phos.
Format	IgG
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion

Produktbeschreibung	Anti-Mouse IgG Antibody generated in sheep detects reactivity to Mouse IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as...
Klonalität	Polyclonal
Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Sheep Serum, Mouse IgG and Mouse Serum.
Formulierung	Liquid (sterile filtered)
Formel	50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN3
Target-Kategorie	Mouse
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:10,000 - 1:50,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500
Anwendungsbeschreibung	Anti-Mouse IgG Alkaline Phosphatase Antibody has been tested by ELISA and dot blot and is assayed against 1.0 ug of Mouse IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code NPP-10 as a substrate for 30 minutes at room temperature.