BIOZOL

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit IgG anti-Sheep IgG (Fc)-Alk. Phos., MinX none

DNA-SEC-183556

Artikelname	Rabbit IgG anti-Sheep IgG (Fc)-Alk. Phos., MinX none
Artikelnummer	DNA-SEC-183556
Hersteller Artikelnummer	SEC-183556
Alternativnummer	DNA-SEC-183556
Hersteller	dianova
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Sheep
Immunogen	Sheep IgG F(c) fragment
Konjugation	Alk. Phos.
Format	IgG
Spezifität	IgG (Fc)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion

Produktbeschreibung	Anti-Sheep IgG F(c) generated in rabbit is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of sheep IgG and o...
Klonalität	Polyclonal
Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Sheep IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Rabbit Serum, Sheep IgG, Sheep IgG F(c) and Sheep Serum. No reaction was observed against Sheep IgG F(ab).
Formulierung	Liquid (sterile filtered)
Formel	50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN3
Target-Kategorie	Sheep
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:2,000 - 1:10,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500
Anwendungsbeschreibung	This product has been assayed against 1.0 ug of Sheep IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code # NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:1,300 to 1:5,400 of the reconstitution concentration is suggested for this product.