**BIOZOL**

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

**E-Mail:** info@biozol.de

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

# Product Datasheet

## Goat Fab anti-Chicken IgG (H+L)-Biotin, MinX none

## DNA-SEC-183918

| | |
|---|---|
| Artikelname | Goat Fab anti-Chicken IgG (H+L)-Biotin, MinX none |
| Artikelnummer | DNA-SEC-183918 |
| Hersteller Artikelnummer | SEC-183918 |
| Alternativnummer | DNA-SEC-183918 |
| Hersteller | dianova |
| Wirt | Goat |
| Kategorie | Antikörper |
| Applikation | ELISA,IHC,WB |
| Spezies Reaktivität | Gallus |
| Immunogen | Chicken IgG whole molecule |
| Konjugation | Biotin |
| Format | Fab |
| Spezifität | IgG (H+L) |
| Minimale Kreuzreaktivität (MinX) | no cross-adsorbtion |

| | |
|---|---|
| Produktbeschreibung | Fab Anti-Chicken IgG Biotin Antibody generated in goat detects chicken IgY. This product possesses the F(ab) region possessing the epitope-recognition site, both heavy and light chains of the antibody molecule are present.... |
| Klonalität | Polyclonal |
| Konzentration | 1.0 mg/mL |
| Isotyp | Ig |
| Puffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Reinheit | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Chicken IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin and anti-Goat Serum. No reaction was observed against anti-Papain or anti-Goat IgG F(c). |
| Formulierung | Lyophilized |
| Formel | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3 |
| Target-Kategorie | Chicken |
| Antibody Type | Secondary Antibody |
| Application Verdünnung | ELISA Dilution: 1:80,000 - 1:800,000, Fluorochrome Protein Value: 10-20, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000 |
| Anwendungsbeschreibung | Suitable for immunoblotting, ELISA, immunohistochemistry, immunomicroscopy as well as other antibody based assays using streptavidin or avidin conjugates requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. |