BIOZOL

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit Fab anti-Goat IgG (H+L)-FITC, MinX none

DNA-SEC-183924

Artikelname	Rabbit Fab anti-Goat IgG (H+L)-FITC, MinX none
Artikelnummer	DNA-SEC-183924
Hersteller Artikelnummer	SEC-183924
Alternativnummer	DNA-SEC-183924
Hersteller	dianova
Wirt	Rabbit
Kategorie	Antikörper
Applikation	FLISA,FACS,IF
Spezies Reaktivität	Goat
Immunogen	Goat IgG whole molecule
Konjugation	FITC
Format	Fab
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion

Produktbeschreibung	Fab Anti-Goat IgG Fluorescein Antibody generated in rabbit detects goat IgG. This product possesses the F(ab) region possessing the epitope-recognition site, both heavy and light chains of the antibody molecule are present. Secondary Antibodies are a...
Klonalität	Polyclonal
Konzentration	0.5 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein and anti-Rabbit Serum. No reaction was observed against anti-Papain or anti-Rabbit IgG F(c).
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target-Kategorie	Goat
Antibody Type	Secondary Antibody
Application Verdünnung	FLISA Dilution: 1:10,000 - 1:50,000, Flow Cytometry Dilution: 1:500 - 1:2,500, Fluorochrome Protein Value: 1.3, IF Microscopy Dilution: 1:1,000 - 1:5,000
Anwendungsbeschreibung	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.