

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Hu CD56 Purified, Clone: [MEM-188], Monoclonal EXB-11-231-C100

| | |
|--------------------------|---|
| Artikelname | Anti-Hu CD56 Purified, Clone: [MEM-188], Monoclonal |
| Artikelnummer | EXB-11-231-C100 |
| Hersteller Artikelnummer | 11-231-C100 |
| Alternativnummer | EXB-11-231-C100 |
| Hersteller | EXBIO |
| Kategorie | Antikörper |
| Applikation | FC, ICC, IHC-P, IP |
| Spezies Reaktivität | Human, Primate |
| Immunogen | KG-1 human acute myelogenous leukemia cell line |
| Produktbeschreibung | CD56 (NCAM, neural cell adhesion molecule) is a transmembrane glycoprotein of immunoglobulin family serving as adhesive molecule which is ubiquitously expressed in nervous system, usually as 120 kDa, 140 kDa or 180 kDa isoform, and it is also found o... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/ml |
| Klon-Bezeichnung | [MEM-188] |
| Isotyp | Mouse IgG2a |
| Puffer | Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| Lagerung | 2°C to 8°C |

| | |
|------------------------|--|
| Target-Kategorie | CD56 |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Flow cytometry: Recommended dilution: 3 µg/ml. Immunohistochemistry (paraffin sections): Recommended dilution: 1:50, pretreatment: heat retrieval (microwave oven) in citrate buffer, standard ABC visualization, positive tissue: membrane-specific reaction in adjacent peripheral nerve, staining of paraprostatic autonomic ganglion. |